

A variational Jastrow coupled-cluster theory of quantum many-body systems

Y. Xian

School of Physics and Astronomy, The University of Manchester, Manchester M13 9PL, UK

(Dated: February 2, 2008)

Abstract

We study many-body correlations in the ground states of a general quantum system of bosons or fermions by including an additional Jastrow function in our recently proposed variational coupled-cluster method. Our approach combines the advantages of state-dependent correlations in the coupled-cluster theory and of strong, short-ranged correlations of the Jastrow function. We apply a generalized linked-cluster expansion for the Jastrow wavefunction and provide detailed analysis for practical evaluation of Hamiltonian expectation value as an energy functional of the Jastrow function and the bare density-distribution functions introduced and calculated in our earlier publications; a simple, first-order energy functional is derived and detailed formulas for higher-order contributions are provided. Our energy functional does not suffer the divergence as in most coupled-cluster calculations when applying to Hamiltonians with hardcore potentials. We also discuss relations between our energy functional and the energy functionals from other theories.

PACS numbers: 03.65.Ca, 31.15.Dv, 31.15.Pf, 71.10.-w, 71.45.-d

I. INTRODUCTION

Most microscopic quantum many-body theories developed over the last five decades can perhaps be broadly divided into two categories, one in real space and the other in momentum (configurational) space. While a real-space many-body theory usually focuses on the interaction potential part of Hamiltonian and evaluates the Hamiltonian expectation value in a first quantization form, a momentum-space theory often starts from the kinetic part of Hamiltonian, is a basis of or closely related to many-body perturbation theories, and mostly deals with Hamiltonian in a second quantization form [1, 2]. A typical real-space approach to the ground state of a quantum many-body system is provided by Jastrow wavefunction which is constructed by a state-independent two-body correlation function [3]. Systematic techniques based on Jastrow wavefunction including extension to inhomogeneous boson or fermion systems is now generally referred to as the method of correlated basis functions (CBF) [4, 5, 6]. The CBF method has proved to be efficient in dealing with strong, short-ranged correlations typified by those in quantum helium liquids [4]. On the other hand, momentum-space many-body theories are often easier to apply and, due to inclusion of the state-dependent correlations, is capable of producing accurate results for a wide range of quantum systems, such as boson gas [7], quantum antiferromagnets with Néel order [8], finite nuclei [9], and electron systems such as electron gas [10], atoms and molecules [11]. A typical momentum theory is the coupled-cluster method (CCM) in which wavefunction are explicitly constructed by state-dependent operators [12, 13]. State-of-the-art calculations of the CCM with high accuracy have often been carried out in quantum chemistry [14], and recently in quantum spin lattices with Néel order [15]. Systematic resummations of diagrams in perturbation theory for boson systems have revealed interesting relations between the two approaches, for example, the hyper-netted chain approximation in the CBF method in fact contains a consistent resummation of both infinite ring and infinite ladder diagrams of momentum-space approach [16, 17]. It appears that real-space and momentum-space approaches complement each other and unification of these two approaches may provide a quantitative description applicable to wider range of quantum many-body systems, including in particular the strongly-correlated fermion systems [18].

We recently extended the CCM to a variational formalism in which bra and ket states are now hermitian to one another [19, 20, 21], contrast to the traditional CCM where they are

not [22]. We introduced the hermitian-conjugate pair of important bare density distribution functions for practical and systematical calculations; the traditional CCM was shown to correspond to a simple linear approximation in one set of distribution functions of our variational coupled-cluster method (VCCM). The well-known momentum approaches such as Bogoliubov theory of boson gas, Anderson's spin-wave theory (SWT), and BCS theory of superconductivity [23], can all be explicitly shown as special low-order approximations in both the ground- and excited-state wavefunctions of the VCCM. We have demonstrated by a detailed application to quantum antiferromagnets with Néel order. Approximations beyond SWT improved results for the ground-state properties [20] and new excitation states have also been obtained [21]. Furthermore, our calculations for the bare density distribution functions can be carried out by diagrammatical techniques similar to those employed by the CBF methods [20]. Hence, a bridge between coupled-cluster theory and Jastrow theory is built. We therefore believe it is a natural next step to combine the two methods for a unified description. This is our main purpose in this article. Krotscheck, Kümmel and Zabolitzky (KKZ) made the first attempt in 1980 for fermion systems [18]. They employed the traditional CCM with only the ket state specified and the Jastrow function is fixed before hand; Hamiltonian eigenequation was used to obtain the ground-state energy and equations for the ket-state coefficients. Here we employ explicit ket and bra states of the VCCM and calculate distribution functions in terms of these ket- and bra-state coefficients; the energy functional is derived in terms of the distribution functions and the Jastrow function.

We focus here on formal development of our approach and organize this article as follows. In Sec. II we introduce our correlated wavefunctions and their distribution functions. The generalized linked-cluster expansion technique is employed for calculations of generating functional in terms of the bare distribution functions. In Sec. III we evaluate Hamiltonian expectation value using Jackson-Feenberg transformation; a first-order energy functional is derived and formulas for higher-order contributions are provided. We conclude in Sec. VI with a summary and discussion on the relations between our approach and other many-body theories.

II. DISTRIBUTION FUNCTIONS IN VARIATIONAL JASTROW COUPLED-CLUSTER THEORY

As proposed earlier [20], we consider a general variational wavefunction by adding a Jastrow correlation operator on top of the Coester state $|\Psi^c\rangle$

$$|\Psi\rangle = e^{S^0/2}|\Psi^c\rangle, \quad (1)$$

and we use the following notation

$$\langle\tilde{\Psi}| = \langle\tilde{\Psi}^c|e^{S^0/2}, \quad (2)$$

for the bra state. In Eqs. (1) and (2), the hermitian operator S^0 is the Jastrow correlation operator, and is given in terms of field operators as [2],

$$S^0 = \frac{1}{2} \int dx_1 dx_2 \psi^\dagger(x_1) \psi^\dagger(x_2) u(x_1, x_2) \psi(x_2) \psi(x_1), \quad (3)$$

where field operators $\psi^\dagger(x)$ and $\psi(x)$ obey the usual boson or fermion commutation relations, $u(x_1, x_2)$ is a local, symmetric function $u(x_1, x_2) = u(x_2, x_1)$, and x are particle coordinates including spin degrees of freedom. We also require that $u(x_1, x_2)$ is bound and short-ranged in real space. We do not include in S^0 any single-body operator as it can be easily absorbed in the Coester states to be defined next. The boson or fermion symmetry is contained in the Coester states by definition. More specifically, the Coester ket state is given by, using the convenient notation invented by Arponen and Bishop [24],

$$|\Psi^c\rangle = e^S|\Phi\rangle, \quad (4)$$

where S is constructed by the so-called configurational creation operators $C_I^\dagger$ which are defined with respect to the model state $|\Phi\rangle$ with the nominal index I labeling multi-particle excitation states from the model state $|\Phi\rangle$,

$$S = \sum_I F_I C_I^\dagger, \quad (5)$$

with F_I often referred to as the correlation coefficients. The Coester bra-state $\langle\tilde{\Psi}^c|$ is given by the hermitian conjugate of the ket state,

$$\langle\tilde{\Psi}^c| = \langle\Phi|e^{\tilde{S}}, \quad \tilde{S} = \sum_I \tilde{F}_I C_I \quad (6)$$

in our VCCM, where C_I are the corresponding configurational destruction operators and $\tilde{F}_I$ are the independent, hermitian conjugates of F_I . In the traditional CCM, however, the bra state is parametrized differently from the ket state and is written as [22],

$$\langle \tilde{\Psi}^c | = \langle \Phi | \tilde{S}' e^{-S}, \quad \tilde{S}' = 1 + \sum_I \tilde{F}'_I C_I. \quad (7)$$

where S is defined as in the ket state of Eq. (5), and $\tilde{F}'_I$ is the bra state coefficients which, in general are not hermitian conjugate of F_I . As discussed in Appendix A, the CCM states violate the condition for application of the generalized linked-cluster expansion due to the *linear* construction of the bra state. Furthermore, the evaluation of the kinetic energy discussed in Appendix B will contain a three-body terms because the nonhermitian relation between the ket and bra states in the CCM. We therefore will not discuss the CCM further and focus only on the VCCM basis of Eqs. (4-6). Clearly, the natural variational parameters are $(F, \tilde{F}, u)$ in the VCCM basis, where we have used the notations $F = \{F_I\}$, $\tilde{F} = \{\tilde{F}_I\}$ and $u = u(x_1, x_2)$. In principle, if these Coester states are exact (namely, all configurations are included in the summations over all I -indices), parameter u is redundant. However, as we always need to make a finite truncation approximation in summations over I -indices in any practical application and it is well-known that the Coester states in a finite truncation approximation are not efficient in dealing with the strong, short-ranged correlations, the two-body Jastrow function $u(x_1, x_2)$ is a useful, important variational parameter in a real application. We hence always assume the summations in Eqs. (5) and (6) are within the subset of a truncation approximation. One of such truncations is the so-called SUB m approximation in which we retain up to m -body creation operators only.

Our basic strategy for calculations is to evaluate the generating functional W of Eqs. (1) and (2),

$$W = \ln \langle \tilde{\Psi} | \Psi \rangle = W^c + W^u, \quad (8)$$

where W^c is the generating function for the pure Coester states without the Jastrow operator,

$$W^c = \ln \langle \tilde{\Psi}^c | \Psi^c \rangle, \quad (9)$$

and W^u is the remainder containing the u function

$$W^u = \ln \langle e^{S^0} \rangle^c, \quad \langle e^{S^0} \rangle^c = \frac{\langle \tilde{\Psi}^c | e^{S^0} | \Psi^c \rangle}{\langle \tilde{\Psi}^c | \Psi^c \rangle}. \quad (10)$$

The general strategy for calculating $W^c = W^c(F, \tilde{F})$ was discussed in our earlier papers and detailed calculations were demonstrated in the spin-lattice application [19, 20]. Briefly, we first introduce the hermitian-conjugate pair of the bare density distribution functions

$$\tilde{g}_I^c = \langle C_I^\dagger \rangle^c = \frac{\partial W^c}{\partial F_I}, \quad g_I^c = \langle C_I \rangle^c = \frac{\partial W^c}{\partial \tilde{F}_I}, \quad (11)$$

where the expectation values $\langle \cdots \rangle^c$ are calculated using the Coester states of Eqs. (4-6) as was given in Eq. (10); any physical quantity is expressed in terms of these distribution functions; these distribution functions are then calculated either by algebraic technique [19] or by diagrammatic technique [20]. In the algebraic approach we derive the following general and self-consistency equations for the calculations,

$$\tilde{g}_I^c = G_I(\tilde{F}, g^c), \quad g_I^c = G_I(F, \tilde{g}^c) \quad (12)$$

where $G_I(\tilde{F}, g^c)$ with notation $g^c = \{g_I^c\}$ etc. as before is a function containing up to linear terms in g^c and finite-order terms in $\tilde{F}$, and similarly for $G_I(F, \tilde{g}^c)$. In the diagrammatic approach, many techniques employed by the CBF method are also applicable here. For example, the coefficients $\tilde{F}_I$ can be replaced in every diagram of W^c expansion by the bare distribution functions $\tilde{g}_I$ after resummations of infinite ring diagrams. In general, the energy expectation value can be expressed as a functional polynomial in F and $\tilde{g}^c$, $E^c = E^c(F, \tilde{g}^c)$; and the variational equations can then be derived by the functional derivatives [25].

In terms of the distribution functions as discussed above, we now consider evaluation of $\langle e^{S^0} \rangle^c$ in Eq. (10). After cluster expansion for the Jastrow part of the wavefunctions, we apply the generalized linked-cluster theorem discussed in Appendix A in a similar fashion as the traditional Jastrow theory [2, 26], and we obtain

$$W^u = \ln \langle e^{S^0} \rangle^c = \sum_{n=2} \frac{1}{n!} \int dx_1 \cdots dx_n [Z_n \rho_n^c]_L \quad (13)$$

where $Z_n(Y)$ are the n -body Yvon-Mayer functions of the bound function Y_{12} ,

$$Y_{12} = e^{u(x_1, x_2)} - 1; \quad (14)$$

ρ_n^c is the n -body density distribution functions of the Coester states; and the notations $[\cdots]_L$ denote that only the linked components of the products are included. In deriving Eq. (13) in Appendix A, we have assumed that the Coester states satisfy the cluster-decomposition properties (i.e., higher-order ρ_n^c are given by sum of products of lower-order ones plus a

nondecomposable core). While this is true in a SUB2 approximation in our spin lattice application and the SUB2 approximation for the Bose gas and BCS superconductors [20, 27], we can not provide a general proof without the details of truncation approximations employed in the Coester states in a real application. Hence by using Eq. (13), we have made a further restriction in the truncation approximations of the Coester states. In the following discussion this further approximation is understood and we will examine this property in real applications beyond the SUB2 approximation. The following analysis for calculating ρ_n^c in terms of F_I and $\tilde{g}_I^c$ clearly show such examination posing no major difficulty.

As all the Yvon-Mayer diagrams in Z_2 and Z_3 are linked, we simply have

$$[Z_2\rho_2^c]_L = Y_{12}\rho_2^c, \quad \rho_2^c(x_1, x_2) = \langle \psi^\dagger(x_1)\psi^\dagger(x_2)\psi(x_2)\psi(x_1) \rangle^c \quad (15)$$

for 2-body cluster contribution and

$$[Z_3\rho_3^c]_L = (Y_{12}Y_{23} + Y_{23}Y_{31} + Y_{31}Y_{12} + Y_{12}Y_{23}Y_{31})\rho_3^c \quad (16)$$

for the 3-body cluster contributions with the 3-body distribution function

$$\rho_3^c = \langle \psi^\dagger(x_1)\psi^\dagger(x_2)\psi^\dagger(x_3)\psi(x_3)\psi(x_2)\psi(x_1) \rangle^c \quad (17)$$

in the Coester states. From the 4-body function $Z_4(Y)$ and onward, however, there are unlinked Yvon-Mayer diagrams which are to be included in the product $[Z_n\rho_n^c]_L$ only after multiplying with the terms of ρ_n^c to form the linked components, and details of which will depend on applications with a truncation approximation in the Coester states. There is clearly a trade-off between the order of linked-cluster expansion and the order of truncation approximation of the Coester states. We hope to get experience in real applications in future. The additional state-dependent correlations of the Coester states are contained firstly in W^c of Eq. (9) and secondly in the density distribution functions ρ_n^c . Importantly, these density distribution functions can be calculated in terms of F and $\tilde{g}^c$, and by using the linearity theorem of the VCCM [21], we can show that all ρ_n^c functions contain only up to linear terms in $\tilde{g}^c$ and finite-order terms in F . As a demonstration, we consider the two-body function ρ_2^c ,

$$\rho_2^c(x_1, x_2) = \langle A_2 \rangle^c = \frac{1}{\langle \tilde{\Psi}^c | \tilde{\Psi}^c \rangle} \langle \tilde{\Psi}^c | A_2 e^S | \Phi \rangle = \frac{1}{\langle \tilde{\Psi}^c | \tilde{\Psi}^c \rangle} \langle \tilde{\Psi}^c | e^S \bar{A}_2 | \Phi \rangle, \quad (18)$$

where $A_2 = \psi^\dagger(x_1)\psi^\dagger(x_2)\psi(x_2)\psi(x_1)$ and $\bar{A}_2$ is

$$\bar{A}_2 = e^{-S} A_2 e^S = e^{-S} \psi^\dagger(x_1)\psi^\dagger(x_2)\psi(x_2)\psi(x_1) e^S = \bar{\psi}^\dagger(x_1)\bar{\psi}^\dagger(x_2)\bar{\psi}(x_2)\bar{\psi}(x_1), \quad (19)$$

with $\bar{\psi}^\dagger(x) = e^{-S}\psi^\dagger(x)e^S$ etc. Using the nested commutation series, we obtain

$$\bar{\psi}^\dagger(x) = \psi^\dagger(x) + [\psi^\dagger(x), S] + \frac{1}{2!}[[\psi^\dagger(x), S], S] + \cdots = \psi^\dagger(x) + [\psi^\dagger(x), S], \quad (20)$$

where the series terminates at first order as S is constructed by creation operator $C_I^\dagger$ only [12, 19]. Evaluation of $\bar{A}_2|\Phi\rangle$ in general leaves only a constant and creation operators acting on $|\Phi\rangle$, namely,

$$\bar{A}_2|\Phi\rangle = \left[X_{2,0}(F; x_1, x_2) + \sum_I X_{2,I}(F; x_1, x_2) C_I^\dagger \right] |\Phi\rangle, \quad (21)$$

$X_{2,0}$ and $X_{2,I}$ are a two-body function containing up to fourth-order terms in F . Therefore, the two-body density function in the Coester states is, using the definition of Eqs. (11)

$$\rho_2^c(x_1, x_2) = X_{2,0}(F; x_1, x_2) + \sum_I X_{2,I}(F; x_1, x_2) \tilde{g}_I^c. \quad (22)$$

In similar fashion, we derive

$$\rho_n^c = X_{n,0} + \sum_I X_{n,I} \tilde{g}_I^c \quad (23)$$

for the n -body density distribution function in the Coester states, where $X_{n,0}$ and $X_{n,I}$ are the n -body functions containing up to $(2n)$ th-order terms in F . Therefore, the linked-cluster contributions of Eq. (13) is written as

$$W^u = \frac{1}{2} \int dx_1 dx_2 Z_2 \left[X_{2,0} + \sum_I X_{2,I} \tilde{g}_I^c \right] + \frac{1}{6} \int dx_1 dx_2 dx_3 Z_3 \left[X_{3,0} + \sum_I X_{3,I} \tilde{g}_I^c \right] + \cdots, \quad (24)$$

where the remainders are the 4-cluster and higher-order contributions, and their calculations will depend on the details of the truncation schemes employed in the Coester states but the general property of finite-order in F and linear in $\tilde{g}^c$ remains.

Before we consider the density-distribution functions of the Jastrow-Coester states of Eqs. (1) and (2), we need to define biased distribution functions as

$$\tilde{g}_I \equiv \frac{1}{\langle \tilde{\Psi} | \Psi \rangle} \langle \tilde{\Psi}^c | e^{S^0} C_I^\dagger | \Psi^c \rangle, \quad g_I \equiv \frac{1}{\langle \tilde{\Psi} | \Psi \rangle} \langle \tilde{\Psi}^c | C_I e^{S^0} | \Psi^c \rangle. \quad (25)$$

They are so called because they are not defined usually as $\langle C_I^\dagger \rangle$ and clearly $\tilde{g}_I \neq \langle C_I^\dagger \rangle$ due to the fact that $C_I^\dagger$ and $S^0/2$ do not commute in general. Similarly, $g_I \neq \langle C_I \rangle$ by our definition. These biased distribution functions can be calculated by the functional derivative of the generating functional of Eq. (8) as

$$\tilde{g}_I = \frac{\partial W}{\partial F_I} = \tilde{g}_I^c + \frac{\partial W^u}{\partial F_I}, \quad (26)$$

where we have used the definition of Eq. (11). The functional derivative in Eq. (26) can be calculated by Eq. (24),

$$\frac{\partial W^u}{\partial F_I} = \frac{1}{2} \int dx_1 dx_2 Z_2 \left\{ \frac{\partial X_{2,0}}{\partial F_I} + \sum_{I'} \left[\frac{\partial X_{2,I'}}{\partial F_I} + X_{2,I'} (\tilde{g}_{I+I'}^c - \tilde{g}_I^c \tilde{g}_{I'}^c) \right] \right\} + \dots, \quad (27)$$

where we have used the fact that

$$\frac{\partial \tilde{g}_{I'}^c}{\partial F_I} = \frac{\partial^2 W^c}{\partial F_I \partial F_{I'}} = \tilde{g}_{I+I'}^c - \tilde{g}_I^c \tilde{g}_{I'}^c, \quad (28)$$

with $\tilde{g}_{I+I'}^c = \langle C_I^\dagger C_{I'}^\dagger \rangle^c = \langle C_{I+I'}^\dagger \rangle^c$. Clearly $\tilde{g}_{I+I'}^c$ are also bare distribution functions.

Using the fact that $S^0/2$ commutes with density operator $\psi^\dagger(x)\psi(x)$, the single-particle density function, $\rho_1(x) = \langle \psi^\dagger(x)\psi(x) \rangle$ can then be calculated as

$$\rho_1(x) = \frac{1}{\langle \tilde{\Psi} | \tilde{\Psi} \rangle} \langle \tilde{\Psi}^c | e^{S^0} \psi^\dagger(x) \psi(x) e^S | \Phi \rangle = \frac{1}{\langle \tilde{\Psi} | \tilde{\Psi} \rangle} \langle \tilde{\Psi}^c | e^{S^0} e^S \bar{\psi}^\dagger(x) \bar{\psi}(x) | \Phi \rangle \quad (29)$$

where the evaluation of $\bar{\psi}^\dagger \bar{\psi} | \Phi \rangle = (X_{1,0} + \sum_I X_{1,I} C_I^\dagger) | \Phi \rangle$ is similar to that in Eqs. (19-23), hence we have, using the definition for the biased distribution function of Eq. (25) and then using Eq. (26),

$$\rho_1(x) = \rho_1^c(x) + \sum_I X_{1,I}(F; x) \frac{\partial W^u}{\partial F_I}, \quad (30)$$

where $\rho_1^c(x)$ is the one-body density-distribution function of the Coester states as in general given by Eq. (23) and the functional derivative $\partial W^u / \partial F_I$ is given by Eq. (27).

The two-body density distribution function of states of Eq. (1) and (2) can be calculated in similar fashion as ρ_1 shown above. We take a more efficient calculation by the functional derivative as

$$\rho_2(x_1, x_2) = \langle \psi^\dagger(x_1) \psi^\dagger(x_2) \psi(x_2) \psi(x_1) \rangle = 2 \frac{\partial W}{\partial u(x_1, x_2)} = 2 \frac{\partial W^u}{\partial u(x_1, x_2)}. \quad (31)$$

The functional derivatives $\partial W^u / \partial u$ only involve the Yvon-Mayor functions Z_n and it is easy to derive from Eq. (13)

$$\rho_2(x_1, x_2) = e^{u(x_1, x_2)} \rho_2^c(x_1, x_2) + \frac{1}{3} \int dx'_1 dx'_2 dx'_3 \frac{\partial Z_3(x'_1, x'_2, x'_3)}{\partial u(x_1, x_2)} \rho_3^c(x'_1, x'_2, x'_3) + \dots, \quad (32)$$

where ρ_n^c are the n -body density-distribution function in the Coester states as given in general by Eq. (23). We can see immediately from Eq. (32) that the short-ranged correlation function $u(x_1, x_2)$ will play an important role for applications to strongly-correlated systems, where the pure Coester states are known to be inefficient.

In summary, we have calculated the density-distribution functions as functionals of $(F, \tilde{g}^c, u)$ where $\tilde{g}^c$ are the bare density distribution functions discussed in our earlier VCCM papers [19, 20]; these calculations are all straightforward up to and including the third-order cluster contributions as these functionals are all polynomials of F and $\tilde{g}^c$, and u enters into these functionals through Yvon-Mayor functions. The fourth- and higher-order cluster contributions will depend on the details of applications with the truncation approximations employed in the Coester states. The first few terms of one-body and two-body functions are given by Eqs. (30) and (32) respectively. We also want to point out that evaluation of $\langle C_I^\dagger \rangle$ or $\langle C_I \rangle$ (or the off-diagonal density functions such as $\langle \psi^\dagger(x)\psi(x') \rangle$) in general are highly nontrivial due to the fact that these operators do not commute with the Jastrow factor $\exp(S^0/2)$. However, these calculations are not needed in the Hamiltonian expectation value to be discussed in the next section; if necessary these calculations can be carried out in an approximation after we have obtained solutions to the variational equations [28]. We will discuss these calculations in future.

III. EVALUATION OF HAMILTONIAN EXPECTATION VALUE

In evaluating a general Hamiltonian expectation value, we first notice that the kinetic part of Hamiltonian in general does not commute with the Jastrow operator $S^0/2$ in our states of Eqs. (1) and (2). In real space, however, the kinetic operator contains only second-order derivatives in particle coordinates. We want to take this advantage by expressing our states in real space for calculations. As shown in Appendix A, the wavefunctions of the Jastrow-Coester states of Eqs. (1) and (2) in real space are given by a product

$$\Psi = \Psi^c \Psi^u, \quad \tilde{\Psi} = \tilde{\Psi}^c \Psi^u \quad (33)$$

where $\Psi^u = e^{U/2} = \exp[\sum_{i<j} u(x_i, x_j)/2]$ is the familiar Jastrow wavefunction and $\Psi^c(x_1, \dots, x_N)$ and $\tilde{\Psi}^c(x_1, \dots, x_N)$ are real-space wavefunctions of the Coester ket- and bra-states respectively. In general, we do not need to know the explicit functional form of Ψ^c and $\tilde{\Psi}^c$ as our calculations involving them are always carried out in a second quantization form as we show below. It is interesting nevertheless to know that in a low-order SUB2 approximation, many-body function Ψ^c is known explicitly as a partial-wave function, the so-called independent pair functions for boson gas [29] or BCS superconductors [30]. The

Coester wavefunctions Ψ^c and $\tilde{\Psi}^c$ obey proper symmetry, namely they are antisymmetric for fermions and symmetric for bosons under the exchange of any pair $x_i \rightleftharpoons x_j$. In the followings, we assume our states of Eqs. (1) and (2) have a fixed particle number N for convenience. It is easy to extend to particle-number nonconserving states as discussed in Appendix A. Our final results are valid for both cases because they are expressed in terms of density distribution functions.

Evaluation of kinetic energy involving Jastrow function is helped by Jackson-Feenberg transformation [2]. We reproduce the transformation as Eq. (B7) in Appendix B for our wavefunctions of Eqs. (33),

$$\int dX \tilde{\Psi} \nabla_i^2 \Psi = \int dX \left[\tilde{\Psi}^c e^U \nabla_i^2 \Psi^c + \frac{1}{4} \tilde{\Psi}^c e^U (\nabla_i^2 U) \Psi^c - \frac{1}{4} e^U \nabla_i^2 (\tilde{\Psi}^c \Psi^c) \right]. \quad (34)$$

Another equivalent expression is also derived as Eq. (B8),

$$\int dX \tilde{\Psi} \nabla_i^2 \Psi = \int dX \tilde{\Psi}^c e^U \left[\nabla_i^2 + \frac{1}{4} (\nabla_i^2 U) + \frac{1}{2} (\nabla_i U) \cdot \nabla_i \right] \Psi^c. \quad (35)$$

Both transformations involve one- and two-body density distribution functions only, and the Jastrow factor e^U appears on the right of the derivatives. The biased distribution functions defined in Eq. (25) are then applicable. The expectation value of a general Hamiltonian with an external field (and/or chemical potential) $A(x)$ is calculated as

$$E = \frac{1}{\langle \tilde{\Psi} | \Psi \rangle} \int dX \tilde{\Psi} \left[-\frac{\hbar^2}{2m} \sum_i \nabla_i^2 + \sum_i A(x_i) + \sum_{i < j} v(x_i, x_j) \right] \Psi. \quad (36)$$

Using Eq. (34), we derive

$$E_1 = \int dx \left[-\frac{\hbar^2}{2m} \rho_1'(x) + \hat{A}_{eff}(x) \rho_1(x) \right] + \frac{1}{2} \int dx_1 dx_2 v_{eff}(x_1, x_2) \rho_2, \quad (37)$$

where ρ_1 and ρ_2 are the one- and two-body density distribution functions; $\hat{A}_{eff}(x)$ is the effective external field operator

$$\hat{A}_{eff} = A(x) - \frac{\hbar^2}{8m} (\nabla^c)^2 \quad (38)$$

with operator ∇^c defined as applying to the Coester states only; v_{eff} is the effective potential defined as

$$v_{eff}(x_1, x_2) = v(x_1, x_2) - \frac{\hbar^2}{8m} (\nabla_1^2 + \nabla_2^2) u(x_1, x_2); \quad (39)$$

and finally ρ'_1 is one-body density function derived from the first term of Eq. (34) and written in the second quantization form as

$$\rho'_1(x) = \frac{1}{\langle \tilde{\Psi} | \Psi \rangle} \langle \tilde{\Psi}^c | e^{S^0} T_1 | \Psi^c \rangle, \quad T_1(x) = \psi^\dagger(x) \nabla^2 \psi(x). \quad (40)$$

Using Eq. (35), we derive second equivalent energy functional as

$$E_2 = \int dx \left[-\frac{\hbar^2}{2m} \rho'_1(x) + A(x) \rho_1(x) \right] + \frac{1}{2} \int dx_1 dx_2 \left[v_{eff}(x_1, x_2) \rho_2 - \frac{\hbar^2}{4m} \rho'_2(u) \right], \quad (41)$$

where $\rho'_2(u)$ is two-body density functions derived from the third integrals of Eq. (35),

$$\rho'_2(u) = \frac{1}{\langle \tilde{\Psi} | \Psi \rangle} \langle \tilde{\Psi}^c | e^{S^0} T_2(u) | \Psi^c \rangle, \quad (42)$$

with the two-body operator $T_2(u)$ given by

$$T_2(u) = \psi^\dagger(x_1) \psi^\dagger(x_2) [\nabla_1 u(x_1, x_2) \cdot \nabla_1 + \nabla_2 u(x_1, x_2) \cdot \nabla_2] \psi(x_2) \psi(x_1). \quad (43)$$

The difference between the two energy functionals is that in E_1 of Eq. (37) we need to take care of the operator ∇^c which applies only to the Coester states and in E_2 of Eq. (41) we need to calculate the two-body density function $\rho'_2(u)$. We hope to get experience in real applications as which form is more practical. The density distribution functions ρ_1 and ρ_2 were calculated earlier by Eqs. (30) and (32). It is easy to show that ρ'_1 and $\rho'_2(u)$ can be calculated in similar fashion. We hence write

$$\rho'_1 = \frac{1}{\langle \tilde{\Psi} | \Psi \rangle} \langle \Psi^c | e^{S^0} T_1 | \Psi^c \rangle = \frac{1}{\langle \tilde{\Psi} | \Psi \rangle} \langle \Psi^c | e^{S^0} e^S \bar{T}_1 | \Phi \rangle \quad (44)$$

where $\bar{T}_1(x) = e^{-S} T_1 e^S = \bar{\psi}^\dagger(x) \nabla^2 \bar{\psi}(x)$. After evaluating $\bar{T}_1(x) | \Phi \rangle = [X'_{1,0}(F; x) + \sum_I X'_{1,I}(F; x) C_I^\dagger] | \Phi \rangle$ as before, we have

$$\rho'_1(x) = \rho_1^c(x) + \sum_I X'_{1,I}(F; x) \frac{\partial W^u}{\partial F_I} \quad (45)$$

where

$$\rho_1^c(x) = \langle T_1 \rangle^c = X'_{1,0}(F; x) + \sum_I X'_{1,I}(F; x) \tilde{g}_I^c \quad (46)$$

is the similar density function of the Coester states, and the derivative $\partial W^u / \partial F_I$ in Eq. (45) is given by Eq. (27). In similar fashion, we derive

$$\rho'_2(u; x_1, x_2) = \rho_2^c + \sum_I X'_{2,I} \frac{\partial W^u}{\partial F_I} \quad (47)$$

where

$$\rho_2^c(u; x_1, x_2) = \langle T_2 \rangle^c = X'_{2,0} + \sum_I X'_{2,I} \tilde{g}_I^c, \quad (48)$$

and $X'_{2,0}$ and $X'_{2,I}$ are obtained by evaluating $\bar{T}_2(u)|\Phi\rangle = (X'_{2,0} + \sum_I X'_{2,I} C_I^\dagger)|\Phi\rangle$.

Eqs. (45) and (47), together with density distributions of Eq. (30) and (32), are all we need for calculating the two equivalent energy functional of Eq. (37) and (41). These are our main results in this paper. Denoting the three terms from the first term of ρ'_1 , ρ_1 and ρ_2 as ϵ , and the higher-order remainders as ΔE_l with $l = 1, 2$ for the two energy functionals, we rewrite the energy functionals of Eqs. (37) and (41) as

$$E_l(F, \tilde{g}^c, u) = \epsilon(F, \tilde{g}^c, u) + \Delta E_l(F, \tilde{g}^c, u), \quad l = 1, 2 \quad (49)$$

where $\epsilon(F, \tilde{g}^c, u)$ is given by,

$$\epsilon = K_1^c + \int dx A(x) \rho_1^c(x) + \frac{1}{2} \int dx_1 dx_2 v_{eff}(x_1, x_2) e^{u(x_1, x_2)} \rho_{12}^c(x_1, x_2), \quad (50)$$

with $K_1^c = -(\hbar^2/2m)\langle T_1 \rangle^c$ for the kinetic energy of the VCCM states. We notice that both energy functions give the same first order approximation ϵ as the term containing $(\nabla^c)^2 \rho_1^c(x)$ in E_1 vanishes after integration by parts. Furthermore, this first-order energy functional ϵ is nothing but the energy functional of the VCCM after replacing the bare potential $v \rightarrow V = v_{eff} e^u$, namely

$$\epsilon(F, \tilde{g}^c, u) = E^c(F, \tilde{g}^c)|_{v \rightarrow V}, \quad V(v, u) = \left[v(x_1, x_2) - \frac{\hbar^2}{8m} (\nabla_1^2 + \nabla_2^2) u(x_1, x_2) \right] e^{u(x_1, x_2)}. \quad (51)$$

This is convenient indeed as no new calculations are needed after the VCCM calculations have been done. We want to emphasize that, in addition to this simple, intuitively appealing approximation of Eq. (51), our main purpose here is to provide the detailed formulas for calculating the higher-order terms in ΔE_l of Eq. (49) in the forms of four equations for ρ_1 , ρ'_1 , ρ_2 and ρ'_2 respectively.

IV. SUMMARY AND DISCUSSION

In summary, we have calculated the Hamiltonian expectation value of a general quantum many-body system as energy functionals of the Coester-state coefficients $\{F_I\}$, bare distribution functions $\{\tilde{g}_I^c\}$ of the Coester states and the Jastrow correlation function $u(x_1, x_2)$. Two

equivalent energy functionals are derived by Eqs. (37) and (41). A single simple, first-order energy from both expressions is derived as the usual VCCM energy functional but with a new potential $V(v, u) = v_{eff}e^u$. The formulas for practical calculations of the higher-order terms are provided in details. It is easy to see that, due to the short-ranged Jastrow factor, our energy functionals do not suffer the divergence as in most coupled-cluster calculations when potential $v(x_1, x_2)$ approaching hardcore potentials. We also want to emphasize that an assumption has been made in applying the generalized linked-cluster expansion of Eq. (13) for the Coester states. We have proved this is valid for a SUB2 truncation in the Coester states of spin lattice application (and similar approximations for Bose gas and BCS state), we need to examine this validity for the Coester states in a specific truncation approximation for a real application. We believe this will not pose major difficulty as we have the simple relations between the full distribution functions and the bare distribution functions of Eqs. (23). We also like to point out that the generalized linked-cluster expansion of Eq. (13) can not be applied to the traditional CCM due to the linear construction of its bra state.

A similar wavefunctions to Eqs. (33) were employed by Owen for study of spin-dependent correlations in nuclear matter [31]. In particular, Ψ^c was approximated by the product of an independent pair function (spin-dependent) and the Slater determinant; and in the cluster expansion, the Jastrow function and the independent pair function are treated together. This differs from our approach. Our calculations involving Coester states are always in the second quantization form and are applicable to higher-order truncation approximations. As mentioned in Introduction, KKZ [18] employed the Coester ket state of Eq. (1), and their energy and the equations for correlation coefficients were obtained by the Hamiltonian eigenequation. The validity of this procedure may be questionable as the Coester ket state was approximated in a truncation approximation. In the traditional CCM approach, the use of Hamiltonian eigenequation is completely equivalent in any truncation approximation to a bi-orthogonal variational equations where the specific bra state of Eq. (7) is employed [22, 24], due to the fact that the normalization integral is always unity and that the bra state is linear in the coefficients. This is certainly not the case after inclusion of the Jastrow function. In particular, the normalization integral in KKZ's equations must contain an additional term involving the bra-state coefficients of Eq. (7). It is not clear how this change will complicate their analysis. However, we agree with KKZ and believe that the Jastrow theory and coupled-cluster theory in principle complement each other and combination of

these two theories may overcome each other's difficulties. The approach presented here is our such attempt and we have taken the advantage of the recent progress in the coupled-cluster theory.

It is interesting to compare our energy functional $E(F, \tilde{g}^c, u)$ with the counterpart in the traditional Jastrow theory, $E(a, u)$, where a is the one-body function. Clearly, the missing state-dependent correlations in $E(a, u)$ are now included in $E(F, \tilde{g}^c, u)$ in terms of F and $\tilde{g}^c$. However, a typical calculation of the Jastrow theory practiced today mostly includes resummation of all cluster terms of the linked cluster expansion by the hyper-netted chain (hnc) approximation for bosons or Fermi-hnc approximation for fermions [5, 6] and one optimization route for the boson system is provided by the pair-phonon analysis (PPA) of Campbell and Feenberg [32]. This is possible as the reference states are single-particle states. In our cluster expansion calculations, the reference state is the Coester states which already contain rich correlations including in particular correct long-ranged correlations such as in the SUB2 approximation. The introduction of the Jastrow function is to provide correct description of strong, short-ranged correlations and we expect such scheme may provide reasonable results even if we include a first few cluster terms. It will also be interesting to investigate the relations between our approach and the PPA of Campbell and Feenberg.

Acknowledgments

This work is part of our long-term research project. It is not possible without the supports and encouragements from colleagues and friends. In particular, this research was initially inspired by discussions with J. Arponen (deceased in 2006) and R.F. Bishop and by their works on an extension of the CCM method [22, 24]; we clearly share a goal to extend the traditional, well-established many-body techniques to a more general formalism capable of more accurate, consistent, complete description and applicable to a wider range of quantum many-body systems.

APPENDIX A: Δ -REPRESENTATION AND GENERALIZED LINKED-CLUSTER EXPANSION

We first consider the conversion relations between real-space wavefunctions and the corresponding states in momentum-space. Let $|x_1, \dots, x_N\rangle$ be the single-particle basis of a N -particle system, namely

$$|x_1, \dots, x_N\rangle = \frac{1}{\sqrt{N!}} \psi^\dagger(x_N) \cdots \psi^\dagger(x_1) |0\rangle \quad (\text{A1})$$

where $\psi^\dagger(x)$ are the field operators and $|0\rangle$ is the corresponding vacuum state. The unity operator is then given by

$$1 = \sum_N \int dX |x_1, \dots, x_N\rangle \langle x_1, \dots, x_N|, \quad (\text{A2})$$

where $\int dX = \int dx_1 \cdots dx_N$. The inner product between two arbitrary states can be written as

$$\langle \Psi_1 | \Psi_2 \rangle = \sum_N \int dX \langle \Psi_1 | x_1, \dots, x_N \rangle \langle x_1, \dots, x_N | \Psi_2 \rangle = \sum_N \int dX \Psi_1^* \Psi_2, \quad (\text{A3})$$

where $\Psi_1 = \Psi_1(x_1, \dots, x_N) = \langle x_1, \dots, x_N | \Psi_1 \rangle$ and $\Psi_2 = \Psi_2(x_1, \dots, x_N) = \langle x_1, \dots, x_N | \Psi_2 \rangle$ are the corresponding wavefunctions in real space with proper boson or fermion symmetry. If these two states have a fixed particle number N , the above inner product is then simply given by the familiar quantum mechanics formula

$$\langle \Psi_1 | \Psi_2 \rangle = \int dX \Psi_1^* \Psi_2, \quad (\text{A4})$$

because $\langle x_1, \dots, x_{N'} | \Psi_2 \rangle = \delta_{N',N} \Psi_2(x_1, \dots, x_N)$. In the followings, we assume our states have a fixed particle number N for convenience. It is easy to extend to particle-number nonconserving states by using Eq. (A3). Our final results in this paper are valid for both cases because they are expressed in terms of density distribution functions.

We denote $\Psi^c(x_1, \dots, x_N) = \langle x_1, \dots, x_N | \Psi^c \rangle$ as the Coester state function in real space, where Coester state $|\Psi^c\rangle$ is given by Eq. (4). The application of the Jastrow operator S^0 of Eq. (3) on the single-particle basis can be written as

$$\frac{S^0}{2} |x_1, \dots, x_N\rangle = \frac{U}{2} |x_1, \dots, x_N\rangle, \quad U = \sum_{i < j} u(x_i, x_j) \quad (\text{A5})$$

or

$$e^{S^0/2} |x_1, \dots, x_N\rangle = e^{U/2} |x_1, \dots, x_N\rangle. \quad (\text{A6})$$

We therefore have, using the fact that $u(x_i, x_j)$ is a real function,

$$\Psi(x_1, \dots, x_N) = \langle x_1, \dots, x_N | e^{S^0/2} | \Psi^c \rangle = e^{U/2} \langle x_1, \dots, x_N | \Psi^c \rangle = \Psi^u \Psi^c, \quad (\text{A7})$$

where $\Psi^u = e^{U/2} = \exp(\sum_{i<j} u(x_i, x_j)/2)$ is the familiar Jastrow wavefunction. Similarly, the bra-state wavefunction is written as

$$\tilde{\Psi}(x_1, \dots, x_N) = \tilde{\Psi}^c \Psi^u = \tilde{\Psi}^c e^{U/2}, \quad (\text{A8})$$

where $\tilde{\Psi}^c$ are the corresponding bra-state wavefunctions of Eqs. (6) or (7) in real space. The Coester wavefunctions Ψ^c and $\tilde{\Psi}^c$ obey proper symmetry, namely they are antisymmetric for fermions and symmetric for bosons under the exchange of any pair $x_i \rightleftharpoons x_j$. We do not need to know their explicit functional forms as our later calculations are always carried out in second quantization form of momentum space.

We follow the similar analysis for the evaluation of Eq. (13) as in the traditional Jastrow theory [2, 16]. After the usual cluster expansion of the Jastrow wavefunction in terms of Yvon-Mayor functions $Z_n(Y)$ with the bound function $Y_{12} = e^{u(x_1, x_2)} - 1$,

$$e^U = 1 + \sum_{i<j} Z_2(x_i, x_j) + \sum_{i<j<k} Z_3(x_i, x_j, x_k) + \dots, \quad (\text{A9})$$

the expectation in Eq. (13) is written as,

$$\langle e^{S^0} \rangle^c = 1 + \frac{1}{I_c} \sum_{n=2} \frac{N!}{(N-n)!n!} \int dx_1 \dots dx_N \tilde{\Psi}^c Z_n(x_1, \dots, x_n) \Psi^c = 1 + \sum_{n=2} \frac{1}{n!} \int dx_1 \dots dx_n Z_n \rho_n^c \quad (\text{A10})$$

where $I_c = \langle \Psi^c | \Psi^c \rangle$ is the normalization integral, the first few Z_n are given by

$$Z_2 = Y_{12}, \quad Z_3 = Y_{12}Y_{23} + Y_{23}Y_{31} + Y_{31}Y_{12} + Y_{12}Y_{23}Y_{31}, \quad (\text{A11})$$

etc., and ρ_n^c is the n -body density distribution functions

$$\rho_n^c = \langle \psi^\dagger(x_1) \dots \psi^\dagger(x_n) \psi(x_n) \dots \psi(x_1) \rangle^c. \quad (\text{A12})$$

Evaluation of ρ_n^c of the Coester states in terms of bare distribution function $\tilde{g}^c$ were discussed in details by Eqs. (18-23) of Sec. II. Here we need to consider their cluster properties in order to apply the linked-cluster theorem. For this purpose, we introduce the so-called Δ -representation discussed as follows. We consider evaluation of n -body distribution functions in the state $|\Psi^d\rangle$ with the following cluster decomposition property: any higher-order ρ_n^d can

be written as products of lower order ρ_m^d with $m = 1, 2, \dots, n-1$, plus a nondecomposable core Δ_n . Let us demonstrate this property in details. We start with the one-body density distribution matrix, also the one-body core by definition,

$$\rho_1^d(x; x') = \langle \psi^\dagger(x) \psi(x') \rangle^d = \Delta_1(x; x'), \quad (\text{A13})$$

where we have used the usual notation $\langle \dots \rangle^d$ for the expectation value in state $|\Psi^d\rangle$. The two-body distribution (diagonal) function, $\rho_2^d = \langle \psi^\dagger(x_1) \psi^\dagger(x_2) \psi(x_2) \psi(x_1) \rangle^d$ is then given by using this decomposition property,

$$\rho_2^d = \Delta_1(x_1; x_1) \Delta_1(x_2; x_2) - \Delta_1(x_1; x_2) \Delta_1(x_2; x_1) + \Delta_2(x_1, x_2; x_1, x_2), \quad (\text{A14})$$

where the first two terms are as given before for the fermions (for bosons, all terms have positive sign) and the Δ_2 is the 2-body nondecomposable core. We can also write Eq. (A14) in a symbolic notation as

$$\rho_2^d = \Delta_1 * \Delta_1 + \Delta_2. \quad (\text{A15})$$

In similar way, the three-body density distribution function is then given by

$$\rho_3^d = \Delta_1 * \Delta_1 * \Delta_1 + \Delta_1 * \Delta_2 + \Delta_3, \quad (\text{A16})$$

where Δ_3 is the nondecomposable core. All terms from the product $\Delta_1 * \Delta_1 * \Delta_1$ of Eq. (A16) were given in details as diagrams in the references quoted earlier. Similar to these diagrams, the new contributions in products $\Delta_1 * \Delta_2$ also include both unlinked terms such as $\Delta_1(x_1; x_1) \Delta_1(x_2, x_3; x_2, x_3)$, and linked terms such as $\Delta_1(x_1; x_3) \Delta_2(x_2, x_3; x_2, x_1)$. The rules for sign and for the symbolic product $(*)$ can also all be defined in similar fashion as before and we will discuss them in details somewhere else. Such decomposition can be carried to higher-order.

Using the cluster decomposition property as discussed above, we have the key ingredient for the linked-cluster expansion, namely, all contributions in the product $Z_n \rho_n^d$ can be represented by clusters of diagrams. If we denote linked diagrams as Γ_A, Γ_B , etc., a contribution in similar cluster expansion as Eq. (A10) for state $|\Psi^d\rangle$ can then be written as

$$\Gamma = (\Gamma_A)^{\nu_A} (\Gamma_B)^{\nu_B} \dots, \quad A \neq B, \quad (\text{A17})$$

with coefficients

$$N(A, B, \dots) = \frac{n!}{(n_A!)^{\nu_A} (n_B!)^{\nu_B} \dots} \quad (\text{A18})$$

giving by the number of distributing the linked part along the n points of the diagram. We therefore have

$$\sum_{\Gamma_n} \frac{1}{n!} \Gamma_n = \sum_{A,B,\dots} \frac{1}{n!} N(A,B,\dots) (\Gamma_A)^{\nu_A} (\Gamma_B)^{\nu_B} \dots = \exp(\Gamma_A/n_A! + \Gamma_B/n_B! + \dots). \quad (\text{A19})$$

Hence we have the following generalized linked-cluster expansion for the state $|\Psi^d\rangle$,

$$\ln\langle e^{S^0}\rangle^d = \sum_{n=2} \frac{1}{n!} \int dx_1 \dots dx_n [Z_n \rho_n^d]_L, \quad (\text{A20})$$

where the notation $[Z_n \rho_n^d]_L$ denotes the contributions to $Z_n \rho_n^d$ by linked diagrams only. Since all terms in Z_2 and Z_3 are linked, the first two terms in Eq. (A20) are independent of the diagram structures of ρ_2^d and ρ_3^d and we write

$$\ln\langle e^{S^0}\rangle^d = \frac{1}{2} \int dx_1 dx_2 Z_2 \rho_2^d + \frac{1}{6} \int dx_1 dx_2 dx_3 Z_3 \rho_3^d + \sum_{n=4} \frac{1}{n!} \int dx_1 \dots dx_n [Z_n \rho_n^d]_L. \quad (\text{A21})$$

From $n = 4$ and onward, we need to know the diagram details of ρ_n^d for calculating their contributions.

In order to apply the generalized linked-cluster expansion of Eq. (A21) to our Coester states, we need to prove that the Coester states satisfy the cluster decomposition property as discussed above. In our earlier VCCM calculation for spin lattices, we have shown indeed the Coester states satisfy such property in a SUB2 approximation employed, where arbitrary order distribution functions can be calculated by the simple functional derivative $\tilde{g}_{i'j'}/\partial F_{ij} = \tilde{g}_{ij} \tilde{g}_{i'j}$ and these bare distribution functions correspond to the density distribution matrices (similar analysis also applied to the SUB2 state for the Bose gas and the BCS superconductors) [20, 27]. We also notice that a similar so-called SUB m truncation approximation in the Δ -representation can also be defined as the approximation retaining up to m core distribution tensors only. It is intuitive to relate the real-space cluster parametrization by core distribution tensor $\{\Delta_n\}$ in the Δ -representation and momentum-space parametrization by $\{F_I, \tilde{g}_I\}$ in the Coester states; the Coester representation provides a practical way to calculate these core tensors. We will not intend to provide a general proof that the Coester states in any truncation approximation will satisfy the cluster decomposition property. We will adopt a practical strategy and apply the the linked-cluster expansion formula of Eq. (A21) to the Coester states in real applications and examine the cluster property in the particular truncation approximation employed. We believe this will not cause a major difficulty as we use the relation between full distribution functions ρ_n^c and the bare distribution functions $\tilde{g}_I^c$ as given by Eqs. (23).

We also like to point out that the traditional CCM states certainly fail the cluster decomposition property due to the linear construction of the bra state of Eq. (7). This can be easily seen as any expectation in the CCM is always linear in the bra state coefficients, contradictory to the cluster decomposition property. Therefore, the generalized linked-cluster expansion of Eq. (A21) can not be applied to the CCM states.

APPENDIX B: JACKSON-FEENBERG TRANSFORMATION

We next consider the expectation value of the kinetic energy operator using wavefunctions of Eqs. (A7) and (A8). We follow the derivation as given in Ref. 2 but keep using the notation $\tilde{\Psi}_C$ as it is different to the ket-state counterpart in the traditional CCM. Applying the nested commutation formula in the following integral

$$I_i \equiv \int dX \tilde{\Psi} \nabla_i^2 \Psi = \int dX \tilde{\Psi}^c e^U \left(e^{-U/2} \nabla_i^2 e^{U/2} \right) \Psi^c, \quad (\text{B1})$$

we have

$$I_i = \int dX \tilde{\Psi}^c e^U \left[\nabla_i^2 + \frac{1}{2}(\nabla_i^2 U) + (\nabla_i U) \cdot \nabla_i + \frac{1}{4}(\nabla_i U)^2 \right] \Psi^c. \quad (\text{B2})$$

We can also apply the same nested commutation formula differently as,

$$I_i = \int dX \tilde{\Psi}^c \left(e^{U/2} \nabla_i e^{-U/2} \right) e^U \left(e^{-U/2} \nabla_i e^{U/2} \right) \Psi^c,$$

and derive another equivalent expression as

$$I_i = \int dX \tilde{\Psi}^c \left[\nabla_i - \frac{1}{2}(\nabla_i U) \right] e^U \left[\nabla_i + \frac{1}{2}(\nabla_i U) \right] \Psi^c. \quad (\text{B3})$$

Summing Eqs. (B2) and (B3), the two $(\nabla_i U)^2$ terms (3-body terms) cancel each other out, we have

$$\begin{aligned} 2I_i = & \int dX \left\{ \tilde{\Psi}^c e^U \nabla_i^2 \Psi^c + \frac{1}{2} \tilde{\Psi}^c e^U (\nabla_i^2 U) \Psi^c + \tilde{\Psi}^c \nabla_i e^U \nabla_i \Psi^c \right. \\ & \left. + \frac{1}{2} \tilde{\Psi}^c \left[e^U (\nabla_i U) \nabla_i + \nabla_i e^U (\nabla_i U) \right] \Psi^c \right\}, \end{aligned}$$

and integrating by parts for the third and last terms, we have

$$2I_i = \int dX \left[\tilde{\Psi}^c e^U \nabla_i^2 \Psi^c + \frac{1}{2} \tilde{\Psi}^c e^U (\nabla_i^2 U) \Psi^c - (\nabla_i \tilde{\Psi}^c) e^U (\nabla_i \Psi^c) \right] + 2B_i, \quad (\text{B4})$$

where we have introduced B_i as

$$B_i \equiv \frac{1}{4} \int dX \left[\tilde{\Psi}^c e^U (\nabla_i U) \cdot (\nabla_i \Psi^c) - e^U (\nabla_i \tilde{\Psi}^c) \cdot (\nabla_i U) \Psi^c \right]. \quad (\text{B5})$$

If $\tilde{\Psi}^c = \Psi^c$, these two terms cancel each other out and $B_i = 0$, we therefore have

$$I_i = \int dX \left[\frac{1}{2} \tilde{\Psi}^c e^U \nabla_i^2 \Psi^c + \frac{1}{4} \tilde{\Psi}^c e^U (\nabla_i^2 U) \Psi^c - \frac{1}{2} (\nabla_i \tilde{\Psi}^c) e^U (\nabla_i \Psi^c) \right]. \quad (\text{B6})$$

Using the identity

$$\nabla_i^2 (\Psi^c)^2 = 2(\nabla_i \Psi^c)^2 + 2\Psi^c \nabla_i^2 \Psi^c,$$

the well-known Jackson-Feenberg formula is then derived as, with $\tilde{\Psi}^c = \Psi^c$,

$$I_i = \int dX \left[\tilde{\Psi}^c e^U \nabla_i^2 \Psi^c + \frac{1}{4} \tilde{\Psi}^c e^U (\nabla_i^2 U) \Psi^c - \frac{1}{4} e^U \nabla_i^2 (\tilde{\Psi}^c \Psi^c) \right]. \quad (\text{B7})$$

Eq. (B6) can also be written as after integrating by part for the last term,

$$I_i = \int dX \tilde{\Psi}^c e^U \left[\nabla_i^2 \Psi^c + \frac{1}{4} (\nabla_i^2 U) \Psi^c + \frac{1}{2} (\nabla_i U) \cdot (\nabla_i \Psi^c) \right]. \quad (\text{B8})$$

We notice that in Eqs. (B7) and (B8) we have assumed the Coester ket- and bra-state wavefunctions equal, $\tilde{\Psi}^c = \Psi^c$. This is true for the VCCM states of Eq. (4-6) but not true in general for the CCM state of Eq. (7) in any finite truncation in S and $\tilde{S}'$. Therefore the two terms in B_i of Eq. (B5) do not cancel in general in the the CCM basis. Our conclusion is that the kinetic expectation value in the CCM basis will also contain three-body density distribution functions.

-
- [1] D.J. Thouless, *The quantum mechanics of many-body systems*, Academic Press, New York, 1961.
 - [2] J.-P. Blaizot and G. Ripka, *Quantum theory of finite systems*, MIT Press, 1986.
 - [3] R. Jastrow, Phys. Rev. **98**, 1479 (1955).
 - [4] J.W. Clark and E. Feenberg, Phys. Rev. **113**, 388 (1959); E. Feenberg, *Theory of quantum fluids*, Academic Press, New York, 1969.
 - [5] S. Fantoni and S. Rosati, Nuovo Cim. A **25**, 593 (1975).
 - [6] E. Krotscheck and M.L. Ristig, Nucl. Phys. A **242**, 389 (1975); E. Krotscheck, Phys. Rev. B **31**, 4267 (1985).
 - [7] N.N. Bogoliubov, J. Phys. (Moscow) **11**, 23 (1947); T.D. Lee, C.N. Yang, Phys. Rev. B **105**, 1119 (1957).
 - [8] P.W. Anderson, Phys. Rev. **86**, 694 (1952); T. Oguchi, *ibid.* **117**, 117 (1960).

- [9] B.D. Day, Rev. Mod. Phys. **39**, 719 (1967); H. Kümmel, K.H. Lührmann, and J.G. Zabolitzky, Phys. Rep. C **36**, 1 (1978).
- [10] R.F. Bishop and K.H. Lührmann, Phys. Rev. B **26**, 5523 (1982); K. Emrich and J.G. Zabolitzky, *ibid.* **30**, 2049 (1984).
- [11] O. Sinanoğlu and K.A. Brueckner, *Three approaches to electron correlations in atoms*, Yale U.P., New Haven, Conn., 1970; J. Paldus, J. Čížek, and I. Shavitt, Phys. Rev. A **5**, 50 (1972).
- [12] F. Coester, Nucl. Phys. **7**, 421 (1958); F. Coester and K. Kummel, Nucl. Phys. **17**, 477 (1960).
- [13] J. Čížek, J. Chem. Phys. **45**, 4256 (1966); Adv. Chem. Phys. **14**, 35 (1969).
- [14] R.J. Bartlett, J. Phys. Chem. **93**, 1697 (1989); Theor. Chim. Acta **80**, 71 (1991) and references therein.
- [15] R.F. Bishop, R.G. Hale, and Y. Xian, Phys. Rev. Lett. **73**, 3157 (1994); C. Zeng, D.J.J. Farnell and R.F. Bishop, J. Stat. Phys. **90**, 327 (1998); D.J.J. Farnell and R.F. Bishop, in *Quantum Magnetism* (ed. U. Schollwöck, J. Richter, D.J.J. Farnell and R.F. Bishop), Lecture Notes in Physics, Vol. **645**, Springer-Verlag, Berlin, Page 307, 2004.
- [16] G. Ripka, Phys. Rep. **56**, 1 (1979); A.D. Jackson, A. Lande, and R.A. Smith, Phys. Rev. Lett. **54**, 1469 (1985).
- [17] R.F. Bishop, in *Condensed matter theories* Vol. 10 (eds. M. Casa, M. de Llano, J. Navrro, and A. Polls), Nova Science Publisher, New York, Page 483, 1995.
- [18] E. Krotscheck, H. Kümmel and J.G. Zabolitzky, Phys. Rev. A **22**, 1243 (1980).
- [19] Y. Xian, in *150 years of Quantum Many-body Theories*, ed. by R.F. Bishop, K.A. Gernoth, and N.R. Walet, World Scientific, Page 107, 2001; Y. Xian, Phys. Rev. B **66**, 184427 (2002).
- [20] Y. Xian, Phys. Rev. B **72**, 224438 (2005).
- [21] Y. Xian, Phys. Rev. B **74**, 212401 (2006); J. Phys.: Condens. Matter **19**, 216221 (2007).
- [22] J. Arponen, Ann. Phys. (N.Y.) **151**, 311 (1983).
- [23] J.R. Schrieffer, *Theory of superconductivity*, Benjamin, Reading, MA., 1964.
- [24] J. Arponen, R.F. Bishop, and E. Pajanne, Phys. Rev. A **36**, 2519 (1987).
- [25] The variational equations in Refs. 20-21 were derived using energy functional $E(F, \tilde{F})$. It is easy to show the equivalence of using energy functional $E(F, \tilde{g}^c)$.
- [26] M. Gaudin, J. Gilespe, and G. Ripka, Nucl. Phys. **176A**, 237 (1971); J.W. Clark, in *Progress in particle and nuclear physics*, ed. by D.H. Wilkinson, Vol. 2, Pergamon, Oxford, Page 89, 1979.

- [27] Y. Xian, unpublished.
- [28] M.L. Ristig, P.M. Lam, and J.W. Clark, Phys. Lett. A **55**, 101 (1975); M.L. Ristig and J.W. Clark, Phys. Rev. B **41**, 8811 (1990).
- [29] S. Fantoni, T.M. Nguyen, A. Sarsa and S.R. Shenoy, Phys. Rev. A **66**, 033604 (2002).
- [30] J. Carlson *et al.*, Phys. Rev. Lett. **91**, 50401 (2003).
- [31] J.C. Owen, Ann. Phys. (N.Y.) **118**, 373 (1979).
- [32] C.E. Campbell and E. Feenberg, Phys. Rev. **188**, 396 (1969).